arXiv:1602.04174v1 [math.AC] 12 Feb 2016

RINGS SATISFYING $*$-PROPERTY

KÜRŞAT HAKAN ORAL, BAYRAM ALI ERSOY, AND ÜNSAL TEKIR

Abstract. In this paper we will investigate commutative rings which have the $*$-property. We say that a ring R satisfy $*-$property if for any family of ideals $\{I_\alpha\}_{\alpha\in S}$ of R in which S is an index set, there exists a finite subset S' of S such that $\sqrt{\bigcap_{\alpha\in S} I_\alpha} = \bigcap_{\alpha\in S'} \sqrt{I_\alpha}$. We will show that any integral domain which satisfy $*-$property is a field. Furthermore, these rings are zero-dimensional. After this we give relations between these rings and Artinian rings.

1. Introduction

Throughout this paper, R denotes a commutative ring with identity and the set of all ideals of R is denoted by $L(R)$.

M. Arapovic gives the following charecterization of imbeddibility of a ring into a 0-dimensional ring: A ring R is embeddable in a zero-dimensional ring if and only if R has a family of primary ideals $\{Q_\lambda\}_{\lambda\in\Lambda}$ such that, $(A1)$ $\bigcap_{\lambda\in\Lambda} Q_\lambda = 0$ and $(A2)$ For each $a \in R$, there is an $n \in \mathbb{N}$ such that for all $\lambda \in \Lambda$, if $a \in \sqrt{Q_\lambda}$ then $a^n \in Q_\lambda$ [2, Theorem 7]. Jim Brewer and Fred Richman [3] give an equivalent condition for the condition (A2) in their paper titled "Subrings of zero-dimensional rings" and characterize the condition (A2). A family $\{Q_\lambda\}_{\lambda\in\Lambda}$ of ideals in a ring R satisfies (A2) if and only if for each (countable) subset $\Gamma \subseteq \Lambda$, $\sqrt{\bigcap_{\lambda\in\Gamma} Q_\lambda} = \bigcap_{\lambda\in\Gamma} \sqrt{Q_\lambda}$. With this motivation if we work this equivalent condition, we see that some rings have further from this property. For an example, if R is an Artinian ring then for any subset $\{I_\alpha\}_{\alpha\in S}$ of $L(R)$,

$$\sqrt{\bigcap_{\alpha\in S} I_\alpha} = \bigcap_{\alpha\in S'} \sqrt{I_\alpha}$$

for some finite subset S' of S [Corollary 1]. This equality is called $*$-property. We call a ring R satisfying $*$-property if any family of ideals of the ring R satisfies $*$-property. After this we give the correspondence between such rings and Artinian rings. Then if we use a result of the paper Subrings of zero-dimensional rings by Jim Brewer and Fred Richman, we get that a ring satisfying $*$-property is a zero-dimensional, in the sense of Krull dimension. Further we give an equivalent

Date: June 16, 2013.

2010 *Mathematics Subject Classification.* Primary 13A15, 16E50; Secondary 13E10.

Key words and phrases. Ideals, von Neumann regular rings, Artinian rings.

This paper is in final form and no version of it will be submitted for publication elsewhere.

condition with rings satisfying $*$-property, Artinian rings, zero-dimensional rings and π-regular rings.

2. Rings satisfying $*$-property

Definition 1. *Let R be a ring and S be an index set. If $\sqrt{\bigcap_{\alpha\in S} I_\alpha} = \bigcap_{\alpha\in S'} \sqrt{I_\alpha}$ (for some finite subset S' of S) is satisfied for any arbitrary family $\{I_\alpha\}_{\alpha\in S}$ of ideals of R, then R is said to satisfy $*$-property.*

It is clear that every finite ring satisfies the $*$-property. For a counterexample it is clear that the ring of integers does not satisfy $*$-property.

Theorem 1. *If R satisfies the d.c.c. for radical ideals then R satisfies $*$-property.*

Proof. Let $\{I_\alpha\}_{\alpha\in\Delta}$ be a family of ideals of R. Now consider $\sqrt{\bigcap_{\alpha\in\Delta} I_\alpha}$ for any arbitrary family $\{I_\alpha\}_{\alpha\in S}$ of ideals of R and Δ index set. Since R satisfies the d.c.c for radical ideals, $\sqrt{\bigcap_{\alpha\in\Delta} I_\alpha} = \sqrt{\bigcap_{i=1}^{n} I_{\alpha_i}} = \bigcap_{i=1}^{n} \sqrt{I_{\alpha_i}}$ for some finite subset $\{\alpha_1, \alpha_2, ..., \alpha_n\}$ of Δ where n is a positive integer. □

Corollary 1. *Every Artinian ring satisfies $*$-property.*

Theorem 2. *Let R be an integral domain. If R satisfies $*$-property, then R is a field.*

Proof. Let $I = \bigcap_{\alpha\in S} I_\alpha$ where I_α's are all nonzero ideals of R for $\alpha \in S$. Then $I \neq (0)$. Indeed, if $I = (0)$, then

$$(0) = \sqrt{I} = \sqrt{\bigcap_{\alpha\in S} I_\alpha}$$

and since R is satisfying $*$-property we get

$$(0) = \sqrt{I} = \sqrt{\bigcap_{\alpha\in S} I_\alpha} = \bigcap_{\beta_j\in S'} \sqrt{I_{\beta_j}}$$

where $S' = \{\beta_1, ..., \beta_n\}$. Since all $I_{\beta_j} \neq (0)$, there exists $0 \neq a_{\beta_j} \in I_{\beta_j}$ for all $\beta_j \in S'$. Therefore,

$$0 \neq a = a_{\beta_1} a_{\beta_2} ... a_{\beta_n} \in \bigcap_{\beta_j\in S'} I_{\beta_j} \subset \bigcap_{\beta_j\in S'} \sqrt{I_{\beta_j}} = \sqrt{I}$$

which is a contradiction. Thus $I \neq (0)$. Choose a nonzero element $x \in I$. Since I is the unique smallest non-zero ideal, we get that $(x) = I = (x^2)$. Thus $x = rx^2$ for some $r \in R$. This shows that x is a unit. Hence R is a field. □

Lemma 1. *Let R be a ring satisfying $*$-property. Then every homomorphic image of R satisfies $*$-property.*

Proof. Let $f : R \to R'$ be a surjective homomorphism, R is satisfying $* - property$ and $\{I'_\alpha\}_{\alpha\in S}$ is an arbitrary family of ideals of R'. Then there exists a family of ideals $\{I_\alpha\}_{\alpha\in S}$ such that $f(I_\alpha) = I'_\alpha$ and $\ker(f) \subseteq I_\alpha$. Then we get,

$$\sqrt{\bigcap_{\alpha\in S} I'_\alpha} = \sqrt{\bigcap_{\alpha\in S} f(I_\alpha)} = \sqrt{f\left(\bigcap_{\alpha\in S} I_\alpha\right)} = f\left(\sqrt{\bigcap_{\alpha\in S} I_\alpha}\right).$$

Since R satisfies $*$-property, there exists a finite subset S' of S such that $\sqrt{\bigcap_{\alpha\in S} I_\alpha} = \bigcap_{\alpha\in S'} \sqrt{I_\alpha}$. And so

$$\sqrt{\bigcap_{\alpha\in S} I'_\alpha} = f\left(\bigcap_{\alpha\in S'} \sqrt{I_\alpha}\right) = \bigcap_{\alpha\in S'} f\left(\sqrt{I_\alpha}\right) = \bigcap_{\alpha\in S'} \left(\sqrt{f(I_\alpha)}\right) = \bigcap_{\alpha\in S'} \sqrt{I'_\alpha}.$$

□

Corollary 2. *Let R be a ring satisfying $*$-property and I be an ideal of R. Then R/I satisfies $*$-property.*

Proposition 1. *Let R and R' be rings satisfying $*$-property. Then $R \times R'$ satisfies $*$-property.*

Proof. It is clear. □

M. Arapovic, in his paper "On the embedding of a commutative ring into a 0-dimensional Commutative ring" gives the following charecterization of imbeddibility of a ring into a 0-dimensional ring [2, Theorem 7].

Theorem 3. *[2, Theorem 7] A ring R is embeddable in a zero-dimensional ring if and only if R has a family of primary ideals $\{Q_\lambda\}_{\lambda\in\Lambda}$ such that*
A1. $\bigcap_{\lambda\in\Lambda} Q_\lambda = 0$ *and*
A2. For each $a \in R$, there is $n \in \mathbb{N}$ such that for all $\lambda \in \Lambda$, if $a \in \sqrt{Q_\lambda}$, then $a^n \in Q_\lambda$.

In the paper "Subrings of zero-dimensional rings", Jim Brewer and Fred Richman give an equivalent condition for $(A2)$ and characterize the condition [3, Theorem 2].

Theorem 4. *[3, Theorem 2] A family $\{Q_\lambda\}_{\lambda\in\Lambda}$ of ideals in a ring R satisfies (A2) if and only if for each (countable) subset $\Gamma \subseteq \Lambda$*

$$\sqrt{\bigcap_{\lambda\in\Gamma} Q_\lambda} = \bigcap_{\lambda\in\Gamma} \sqrt{Q_\lambda}\ .$$

Theorem 5. *[3, Theorem 4] The following conditions on a ring R are equivalent:*
(i) The ring R is zero dimensional,
(ii) Condition (A2) holds for the family of all ideals of R,
(iii) Condition (A2) holds for the family of all primary ideals of R.

With this Theorem we give the following Proposition, which gives us that a ring satisfying $*$-property satisfies the condition (A2). And so we get that every ring satisfying $*$-property is a zero-dimensional ring.

Proposition 2. *Let R be a ring satisfying $*$-property and $\{Q_\alpha\}_{\alpha\in S}$ be a family of ideals. Then $\{Q_\alpha\}_{\alpha\in S}$ satisfies the (A2) condition.*

Proof. By assumption for any countable subset $T \subseteq S$,

$$\sqrt{\bigcap_{\alpha\in T} Q_\alpha} = \bigcap_{\alpha\in T'} \sqrt{Q_\alpha}$$

for some finite subset T' of T. Then

$$\bigcap_{\alpha\in T'} \sqrt{Q_\alpha} = \sqrt{\bigcap_{\alpha\in T} Q_\alpha} \subseteq \bigcap_{\alpha\in T} \sqrt{Q_\alpha} \subseteq \bigcap_{\alpha\in T'} \sqrt{Q_\alpha}$$

and so we get

$$\sqrt{\bigcap_{\alpha\in T} Q_\alpha} = \bigcap_{\alpha\in T} \sqrt{Q_\alpha}.$$

Thus by Theorem 4 we get the result. □

Corollary 3. *Every ring which satisfies $*$-property is zero-dimensional.*

Proof. It follows from Proposition 2 and Theorem 5. □

Here we recall that a commutative ring with identity is called von Neumann regular ring (VNR) if for all $x \in R$ there exists $y \in R$ such that $x^2y = x$. Also we give the following characterization of VNR which was given by R.Gilmer in [4].

Theorem 6. *[4, Theorem 3.1] The following conditions are equivalent in a ring R:*
(i) R is VNR,
(ii) R is zero-dimensional and reduced,
(iii) R_P is a field for each $P \in Spec(R)$,
(iv) Each ideal of R is a radical ideal,
(v) Each ideal of R is idempotent.

Corollary 4. *Let R be a reduced ring. If R is satisfying $*$-property then R is VNR.*

Proof. It follows from Corollory 3 and Theorem 6. □

Lemma 2. *Let R be a ring satisfying $*$-property. Consider the following conditions:*
(i) R is a VNR,
(ii) If $\sqrt{I} = \sqrt{J}$ for some ideals I, J of R then $I = J$.
If one of the above conditions holds, then R is an Artinian ring.

Proof. Let R be a ring satisfying $*$-property and

$$I_1 \supseteq I_2 \supseteq I_3 \supseteq ...$$

be a descending chain of ideals in R. Then we get $\sqrt{\bigcap_{\alpha\in S} I_\alpha} = \bigcap_{i=1}^{n} \sqrt{I_{\alpha_i}}$ for some finite subset $S' = \{\alpha_1, ..., \alpha_n\}$ of S, where $I_{\alpha_1} \supseteq ... \supseteq I_{\alpha_n}$. And so

$$\sqrt{\bigcap_{\alpha\in S} I_\alpha} = \bigcap_{i=1}^{n} \sqrt{I_{\alpha_i}} = \sqrt{\bigcap_{i=1}^{n} I_{\alpha_i}} = \sqrt{I_{\alpha_n}}.$$

Thus by our hypothesis we get $\bigcap_{\alpha\in S} I_\alpha = I_{\alpha_n}$. The chain must stop. □

Recall that a commutative ring with identity R is called Laskerian if each ideal of R is a finite intersection of primary ideals.

Theorem 7. *Let R be a commutative ring with identity. If R is a Laskerian ring then the following conditions are equivalent:*
(i) R is a ring satisfying $*$-property,
(ii) For the family $\{P_\lambda\}_{\lambda\in\Lambda}$ of prime ideals of R we get $\sqrt{\bigcap_{\lambda\in\Lambda} P_\lambda} = \bigcap_{\lambda\in\Gamma} P_\lambda$ for a finite subset Γ of Λ .

Proof. (i)⇒(ii) It is clear.
(ii)⇒(i) Suppose that $\sqrt{\bigcap_{\lambda\in\Lambda} P_\lambda} = \bigcap_{\lambda\in\Gamma} P_\lambda$ holds for a finite subset Γ of Λ. Let $\{I_\alpha\}_{\alpha\in\Delta}$ be a family of ideals of R. Since R is Laskerian, we have $\sqrt{I_\alpha} = \bigcap_{i=1}^{n} P_{\alpha,i}$. Thus we get $\sqrt{\bigcap_{\alpha,i} P_{\alpha,i}} = \bigcap_{finite} \sqrt{P_{\alpha,i}} = \bigcap_{finite} P_{\alpha,i}$ and so, $\bigcap_{finite} \sqrt{I_\alpha} = \bigcap_{finite} P_{\alpha,i} = \bigcap_{finite} \sqrt{P_{\alpha,i}} = \sqrt{\bigcap_{\alpha,i} P_\alpha} \subseteq \sqrt{\bigcap_{\alpha} I_\alpha}$. Since the converse inclusion holds always, we get $\sqrt{\bigcap_{\alpha} I_\alpha} = \bigcap_{finite} \sqrt{I_\alpha}$. Hence R satisfies $*$-property. □

Corollary 5. *Let R be a Laskerian ring which satisfies $*$-property and S be a multiplicatively closed subset of R. Then $S^{-1}R$ satisfies $*$-property.*

Proof. Let $\{\mathfrak{P}_\alpha\}$ be a family of prime ideals of $S^{-1}R$. Then there exists a family of prime ideals $\{P_\alpha\}$ of R such that $(P_\alpha)^e = \mathfrak{P}_\alpha$. And we have $\bigcap (P_\alpha)^e = \left(\bigcap P_\alpha\right)^e$ since P_α are prime ideals of R. Thus we get

$$\begin{aligned}\sqrt{\bigcap_{\lambda\in\Lambda} \mathfrak{P}_\lambda} &= \sqrt{\bigcap_{\lambda\in\Lambda} (P_\lambda)^e} = \sqrt{\left(\bigcap_{\lambda\in\Lambda} P_\lambda\right)^e} = \left(\sqrt{\bigcap_{\lambda\in\Lambda} P_\lambda}\right)^e \\ &= \left(\bigcap_{\lambda\in\Gamma} \sqrt{P_\lambda}\right)^e = \left(\bigcap_{\lambda\in\Gamma} P_\lambda\right)^e = \bigcap_{\lambda\in\Gamma} (P_\lambda)^e \\ &= \bigcap_{\lambda\in\Gamma} \mathfrak{P}_\lambda = \bigcap_{\lambda\in\Gamma} \sqrt{\mathfrak{P}_\lambda}\end{aligned}$$

for a finite subset Γ of Λ . Thus $S^{-1}R$ satisfies $*$-property. □

Now, if we let a Noetherian ring instead of Laskerian ring then we can give the following theorem from the previous results. Before the main theorem we will recall that a ring R is called π-regular provided for all $a \in R$ there exist $a' \in R$, $n \in \mathbb{N}^*$ such that $a^n = (a^n)^2 a'$. In his paper "Characterization of the 0-dimensional rings", Arapovic has show that a ring is π-regular if and only if the ring is zero dimensional [1, Theorem 6].

Theorem 8. *Let R be a Noetherian ring. The following conditions are equivalent:*
(i) R satisfies $*$-property,
(ii) R is zero-dimensional,
(iii) R is Artinian,
(iv) R is π-regular,
(v) For the family $\{P_\lambda\}_{\lambda\in\Lambda}$ of prime ideals of R we get $\sqrt{\bigcap_{\lambda\in\Lambda} P_\lambda} = \bigcap_{\lambda\in\Gamma} P_\lambda$ for a finite subset Γ of Λ .

Proof. $(i) \Rightarrow (ii)$ follows from Corollary 3,
$(ii) \Rightarrow (iii)$ follows from [5, Proposition 8.38],
$(iii) \Rightarrow (i)$ follows from Corollary 1,
$(ii) \Leftrightarrow (iv)$ follows from [1, Theorem 6],
$(i) \Leftrightarrow (v)$ follows from Theorem 7. □

References

[1] Arapovic, M., Characterization of the 0-dimensional rings, Glasnik Mathematicki, Vol. 18 (38), 1983, 39-46.
[2] Arapovic, M., On the embedding of a commutative ring into a 0-dimensional commutative ring, Glasnik Mathematicki, Vol. 18 (38), 1983, 53-59.
[3] Brewer, J. and Richman, F., Subrings of zero-dimensional rings, Multiplicative Ideal Theory in Commutative Rings, Springer, 2006, 73-88.
[4] Gilmer, R., Backhround and preliminaires on zero-dimensional rings, Zero Dimensional Rings, Lecture Notes in Pure and Applied Mathematics, 171, Marcel Dekker 1995, 1-13.
[5] Sharp, R. Y., Steps in Commutative Algebra, Cambridge University Press, Second edition 2000.

Yildiz Technical University,Department of mathematics Davutpaşa,34210,Esenler,Istanbul,TURKEY
E-mail address: khoral@yildiz.edu.tr

Current address: Yıldız Technical University,Department of mathematics Davutpaşa,34210,Esenler,Istanbul,TURKEY
E-mail address: ersoya@yildiz.edu.tr

Marmara University, Department of mathematics, Ziverbey, 34722,Goztepe,Istanbul,TURKEY
E-mail address: utekir@marmara.edu.tr